

ABOUT THE Q^2 DEPENDENCE OF THE ASYMMETRY A_1 FROM THE SIMILARITY OF THE g_1 AND F_3 STRUCTURE FUNCTIONS

A.V. KOTIKOV^{1†} and D.V. PESHEKHONOV^{2†}

(1) Bogoliubov Laboratory of Theoretical Physics, JINR, Dubna, 141980 Russia

(2) Laboratory of Particle Physics, JINR, Dubna, 141980 Russia

† E-mail: kotikov@thsun1.jinr.ru, peshehon@sunse.jinr.ru

Abstract

We consider a new approach for taking into account the Q^2 dependence of asymmetry A_1 . This approach is based on the similarity of the Q^2 behavior and the shape of the spin-dependent structure function $g_1(x, Q^2)$ and spin averaged structure function $F_3(x, Q^2)$.

An experimental study of the nucleon spin structure is realized [1] by measuring of the asymmetry $A_1(x, Q^2) = g_1(x, Q^2)/F_1(x, Q^2)$. The most known theoretical predictions on spin dependent structure function (SF) $g_1(x, Q^2)$ of the nucleon were done by Bjorken [2] and Ellis and Jaffe [3] for the so called *first moment value* $\Gamma_1 = \int_0^1 g_1(x) dx$.

Studying the properties of $g_1(x, Q^2)$ and the calculation of the Γ_1 value require the knowledge of SF function g_1 at the same Q^2 in the hole x range. Experimentally asymmetry A_1 is measuring at different values of Q^2 for different x bins. An accuracy of the modern experiments [4]-[7] allows to analyze data in the assumption [8] that asymmetry $A_1(x)$ is Q^2 independent (i.e. SF g_1 and F_1 have the same Q^2 dependence)




Figure 1: The contributions to the structure function g_1^n of the neutron from the valence quarks and from the sea quarks and gluons. The shaded and hatched areas represent the total uncertainties on each quantity. [5] and [7], respectively).

$$A_1(x, Q^2) = A_1(x) \quad (1)$$

But the precise checking of the Bjorken and Ellis - Jaffe sum rules requires considering the Q^2 dependence of A_1 or g_1 . Moreover, the assumption (1) does not follow from the theory. On the contrary, the behavior of F_1 and g_1 as a functions of Q^2 is expected to be different due to the difference between polarized and unpolarized splitting functions.

There are several approaches (see [9] and references therein) how to take into account the Q^2 dependence of A_1 . They are based on different approximate solutions of the DGLAP equations. Some of them have been used already by Spin Muon Collaboration (SMC) and E154 Collaboration in the last analyses of experimental data (see




Figure 2: The structure function $xg_1^p(x, Q^2)$ evolved to $Q^2 = 10 \text{ GeV}^2$ using our Eq.(2), the assumption that g_1^p/F_1^p is Q^2 independent, and exact DGLAP next-to-leading evolution.

and differ from F_1 one already in the first subleading order. For the singlet (S) part of g_1 and for SF F_3 the difference between splitting functions is also negligible (see [11]). This observation allows us to conclude that the ratio of the SF g_1 and F_3 should be essentially Q^2 independent, i.e.

$$A_1^*(x, Q^2) = \frac{g_1(x, Q^2)}{F_3(x, Q^2)} = A_1^*(x) \quad (2)$$

and the asymmetry A_1 at some Q^2 can be defined than as :

$$A_1(x_i, Q^2) = \frac{F_3(x_i, Q^2)}{F_3(x_i, Q_i^2)} \cdot \frac{F_1(x_i, Q_i^2)}{F_1(x_i, Q^2)} \cdot A_1(x_i, Q_i^2), \quad (3)$$

where $x_i (Q_i^2)$ means an experimentally measured value of $x (Q^2)$.

In principle, Q^2 independence of the $A_1^*(x)$ ratio can be violated at small x values, because the SF $F_3(x)$ has only nonsinglet structure and the SF $g_1(x)$ contains gluon and sea quark contributions. We note, however, that

- in polarized case the gluon and sea quark contributions are not so large even at modern small x values (see Fig. 1 which is taken from [7]): only at $x \leq 10^{-3}$ these contributions start to be dominant.
- The double-logarithmic estimations of small x asymptotics for the SF g_1 and F_3 $g_1^{NS}(x)$, $F_3(x) \sim x^{-a_{NS}}$ and $g_1^S(x) \sim x^{-a_S}$, given in [12], lead to the results: $a_{NS} \sim 0.4$ and $a_S \sim 3/2a_{NS} \sim 0.6$, that supports a similarity of nonsinglet and singlet polarized components.

We note that the estimations in [12] have been given at small Q^2 values. However, as it was shown earlier in [13] and [14], respectively, the values of a_{NS} and a_S should be nearly Q^2 -independent, that is supported also by recent fits (see [15] and references therein).

To apply the proposed approach we use the SMC [5] data (the SLAC [6, 7] data have been considered in [11]). We parameterize CCFR data on $F_2(x, Q^2)$ and $xF_3(x, Q^2)$ [16]

In this paper we review another idea which is based on the observation in [10, 11] that the splitting functions of the DGLAP equations for the g_1 and F_3 and the SF shapes are similar in a wide x range and, thus, their Q^2 dependences have to be close as a consequence. Our approach for Q^2 -dependence of A_1 is very simple (see Eq.(3)) and leads to the results, which are very similar to ones based on the DGLAP evolution.

To demonstrate the validity of the observation, we note that splitting functions in the r.h.s. of DGLAP equations for nonsinglet (NS) parts of g_1 and F_3 are the same (at least in first two orders of the perturbative QCD)

and take also the SLAC parameterization of $R(x, Q^2)$ [18] to obtain structure function $F_1(x, Q^2)$. We use in Eq.(3) parameterizations of CCFR data [16] for both SF $xF_3(x, Q^2)$ and $F_2(x, Q^2)$ to avoid systematic uncertainties and nucleon correlation in nuclei.

The SF $g_1(x, Q^2)$ is calculated using the asymmetry $A_1(Q^2)$ as

$$g_1(x, Q^2) = A_1(x, Q^2) \cdot F_1(x, Q^2), \quad (4)$$

where the spin average SF F_1 has been calculated using NMC parameterization of $F_2(x, Q^2)$ [17]. The results are presented in Fig. 2. Our results are in very good agreement with the calculations based on direct DGLAP evolution.

Q^2 (GeV ²)	100	30	10	3
SMC proton and deuteron data				
A_1 -scaling	0.297	0.226	0.202	0.170
Evolution			0.183	
A_1^* -scaling	0.210	0.201	0.191	0.176
SMC proton and E154 neutron data				
A_1 -scaling	0.221	0.209	0.194	0.170
A_1^* -scaling	0.194	0.190	0.185	0.175
Theory	0.194	0.191	0.187	0.180

Table: The mean values of $\Gamma_1^p - \Gamma_1^n$.

the errors of original analyses [5]-[7]. The value of $\Gamma_1^p - \Gamma_1^n$ at $Q^2 = 10$ GeV² obtained by direct DGLAP evolution are taken from article [5].

To make another comparison with the theory we have calculated also the first moment value of the structure function g_1 at different Q^2 . Using Eq.(3), we recalculate the SMC measured asymmetry of the proton and deuteron and E154 one of neutron at $Q^2 = 100$ GeV², 30 GeV², $Q^2 = 10$ GeV² and 3 GeV² and get the value of $\int g_1(x)dx$ through the measured x ranges. To obtain the first moment values $\Gamma_1^{p(d)}$ we use an original estimations of SMC and E154 for unmeasured regions. In Table we present the results for the mean values of $\Gamma_1^p - \Gamma_1^n$, because the errors coincide with

Let us now present the main results, following from the Table and the Fig. 2.

- The results are in very good agreement with $g_1(x, Q^2)$ data of SMC, based on direct DGLAP evolution. So, the suggestion about Q^2 independence of the A_1^* ratio leads to correct results in simplest way.
- Our method allows to test of the Bjorken sum rule in a simple way with a good accuracy. Obtained results on the $\Gamma_1^p - \Gamma_1^n$ show that used experimental data well confirm the Bjorken sum rule prediction.
- A violation of A_1^* -scaling (2) in (future) experimental data will demonstrate large gluon and/or sea quark contributions to SF $g_1(x, Q^2)$. Then, the check of Q^2 dependence of A_1^* ratio should be very useful for an indication of large contributions of gluons and sea quarks in future polarized experiments.

Acknowledgments. One of us (AVK) is grateful very much to Organizing Committee of Workshop DUBNA-SPIN01 for the financial support. A.V.K. was supported in part by Alexander von Humboldt fellowship and INTAS grant N366.

References

- [1] M. Anselmino, A. Efremov and E. Leader, Phys. Rep. **261**, 1 (1995).

- [2] J.D. Bjorken, Phys. Rev. **148**, 1467 (1966); **D1**, 1376 (1970).
- [3] J. Ellis and R.L. Jaffe, Phys. Rev. **D9**, 1444 (1974); **D10**, 1669 (1974).
- [4] SM Collab., B. Adams *et al.*, Phys. Lett. **B329**, 399 (1994); **B357**, 248 (1995); SLAC-E143 Collab., K. Abe *et al.*, Phys. Rev. Lett. **74**, 346 (1995); **75**, 25 (1995); Phys. Rev. **D58**, 112003 (1998).
- [5] SM Collab., B. Adams *et al.*, Phys. Rev. **D56**, 5330 (1997); Phys. Lett. **B396**, 338 (1997).
- [6] SLAC-E154 Collab., K. Abe *et al.*, Phys. Rev. Lett. **97**, 26 (1997).
- [7] SLAC-E154 Collab., K. Abe *et al.*, Phys. Lett. **B405**, 180 (1997).
- [8] J. Ellis and M. Karliner, Phys. Lett. **B313**, 131 (1993); F.E. Close and R.G. Roberts, Phys. Lett. **B316**, 165 (1993).
- [9] R.D. Ball *et al.*, Phys. Lett. **B378**, 255 (1996); M. Gluck *et al.*, Phys. Rev. **D53**, 4775 (1996); T. Gehrmann and W.J. Stirling, Phys. Rev. **D53**, 6100 (1996); A.V. Kotikov and D.V. Peshekhonov, hep-ph/9604269; Phys. Atom. Nucl. **60**, 653 (1997); Phys. Rev. **D54**, 3162 (1996); E. Leader *et al.*, Phys. Lett. **B462**, 189 (1999); S. Simula *et al.*, Preprint RM3-TH/01-4 (hep-ph/0107036).
- [10] A. V. Kotikov and D. V. Peshekhonov, JETP Lett. **65**, 7 (1997); *in* Proc. of the Workshop DIS96 (1996), p.612 (hep-ph/9608369).
- [11] A. V. Kotikov and D. V. Peshekhonov, Eur. Phys. J. **C9**, 55 (1999); *in* the Proc. of the Workshop DIS98 (1998), p.242 (hep-ph/9805374).
- [12] J. Bartels *et al.*, Z. Phys. **C72**, 627 (1996); **C70**, 273 (1996); B.I. Ermolaev *et al.*, Z. Phys. **C69**, 259 (1996); Nucl. Phys. **B594**, 71 (2001); **B571**, 137 (2001); *in* Proc. of the Workshop DIS2001 (2001), (hep-ph/0106317).
- [13] F. Martin, Phys. Rev. **D19**, 1382 (1979); C. Lopez and F.I. Yndurain, Nucl. Phys. **B171**, 231 (1980); A.V. Kotikov *et al.*, Theor. Math. Phys. **84**, 744 (1991).
- [14] L. L. Jenkovszky *et al.*, Sov. J. Nucl. Phys. **55**, 1224 (1992); A.V. Kotikov, Phys. Atom. Nucl. **56**, 1276 (1993); **57**, 133 (1994); **59**, 2137 (1996); Phys. Rev. **D49**, 5746 (1994); Mod. Phys. Lett. **A11**, 103 (1996); A.V. Kotikov *et al.*, Theor. Math. Phys. **111**, 442 (1997).
- [15] A.L. Kataev *et al.*, Preprint CERN-TH/2001-58 (hep-ph/0106221); V.G. Krivokhijine and A.V. Kotikov, Dubna preprint E2-2001-190 (hep-ph/0108224).
- [16] CCFR/NuTeV Collab., W. Seligman *et al.*, Phys. Rev. Lett. **79**, 1213 (1997); W. Seligman, PhD Thesis (Columbia University), Nevis Report 292.
- [17] NM Collab., M. Arneodo *et al.*, Phys. Lett. **B364**, 107 (1995).
- [18] L. W. Whitlow *et al.*, Phys. Lett. **B250**, 93 (1990).